

Hassan Zamani (MD)^{*1}
Kazem Babazadeh (MD)¹
Rahman Ghaffari (MD)²
Hossein Karami (MD)³
Saeid Fattahi (MD)⁴
Farzad Mokhtari Esbuie (MD)⁴

1. Non-Communicable Pediatric Diseases Research Center, Babol University of Medical Sciences, Babol, Iran.
2. Department of Cardiac Surgery, Mazandaran University of Medical Sciences, Sari, Iran.
3. Department Pediatric Oncology, Mazandaran University of Medical Sciences, Sari, Iran.
4. Mazandaran University of Medical Sciences, Sari, Iran.

*** Correspondence:**

Hassan Zamani, Non-Communicable Pediatric Diseases Research Center, No 19, Amirkola Children's Hospital, Amirkola, Babol, Iran.
Postal code: 47317-41151

E-mail: zamanihassan@yahoo.com
Tel: 0098 111 3242151-5
Fax: 0098 111 3246963

Received: 30 Jan 2011
Revised: 31 July 2011
Accepted: 14 Dec 2011

The successful withdrawal of a migrated central venous catheter

Abstract

Background: Central venous catheters (CVCs) have been used widely in clinics. These catheters are also recommended for children and infants receiving chemotherapy and total parenteral nutrition (TPN) and etc. In this paper, we present migrated fractured control line of the heart of a girl.

Case Presentation: A 2.5 year old girl with migrated of the fractured central line into the heart. In the catheterization laboratory, first we placed a long sheath (8 F) into the inferior vena cava via femoral vein and then trapped the foreign body by pigtail catheter and wire 0.035 inch and pulled it down to make its proximal free. After that, we snared the catheter by snare-catheter and pulled it into the femoral vein, and then the cardiac surgeon bridged it out by cut-down successfully.

Conclusion: A rare complication in the use of central catheters is fraction and cardiac embolization. We offer gentle bringing out of the catheter lines under fluoroscopy guide in all of the cases, if this is technically possible and safe.

Keywords: Central Venous Catheter, Foreign body, Cardiac embolization, Snare

Caspian J Intern Med 2012; 3(4): 550-553

Central venous catheters (CVCs) have been used widely in clinics. These catheters are recommended for children and infants receiving chemotherapy and total parenteral nutrition (TPN) and permit the safe infusion of hypertonic solutions and vasoactive agents that can produce severe peripheral soft tissue damage if they extravasate from peripheral line into local tissue. Rapid infusions of large volumes of fluids or blood products are also possible (1). CVCs may be inserted from different sites; including the femoral, subclavian, external jugular, internal jugular, antecubital and, rarely, saphenous veins. The line should terminate at the atriocaval junction to minimize cardiac tamponade. Immediate complications of CVCs include dysrhythmias, pneumothorax, hydrothorax, hemothorax, air embolism, shearing of the vessel, bleeding, apnea, oversedation, and airway obstruction. Central line migration has been described in less than 1% of the patients having central lines. Other complications including fracture and migration or infection may have a higher incidence (2-6). In this article, we present migrated fractured control line of the heart of a girl.

Case presentation

A 2.5 year old girl with acute lymphoblastic leukemia (ALL) had a CVC inserted in subclavian vein previously for chemotherapy, in Boali Hospital, Sari, Iran. The CVC broke during takedown after finishing the course of chemotherapy and then migrated into the heart. In the first chest radiography (P.A view) and fluoroscopy, it was kinked in PA with the proximal and distal tips in the hepatic veins and right ventricular outflow tract (RVOT) (figure 1, 2).

There is complicity of the case because, both tips of the catheter were out of reach and we could not snare them. In the catheterization laboratory, first via femoral vein approach, we placed a long sheath (8 F) into the inferior vena cava and trapped of the foreign body by pigtail catheter and wire 0.035 inch and pulled it down to make its proximal site free. After that, we snared the catheter by snare-catheter and pulled it into the femoral vein, and then the cardiac surgeon brought it out by cutting -down successfully (figure 3, 4).




Figure 1: The localization of proximal and distal tip of catheter in the fluoroscopy




Figure 2: The localization of proximal and distal tip of catheter after the injection of angiography dye contrast.



Figure 3: The snaring of the catheter.




Figure 4: The removed catheter

Discussion

In appropriate manipulation due to a lack of experience in basic techniques of inserting and removing a catheter or cutting a catheter by mistake during removal may result in retention of residual fragments inside the body (7-11) A first case of a polyethylene catheter as a foreign body found in the right atrium at autopsy was reported in 1954 (12). Since then, there have been many reports of unretrieved intravascular devices. The first case of a broken guide wire was reported in 1962, and then reports of similar incidents increased with the introduction of advanced catheter intervention procedures (11, 13-16).

In 1964, the first successful percutaneous removal of a broken guide wire from the right atrium using bronchoscopes forceps was reported, and some similar reports followed (13, 15-17). Nowadays, a variety of catheter devices, including the loop snare catheter, basket catheter, and grasping/biopsy forceps, are developed. By using these devices, foreign bodies could be retrieved safely and promptly.

Loop snare catheters such as the goose-neck snare and Curry's loop snare are widely used (18-20). Loop snare catheters have the advantage of being flexible, allowing them to follow the intravascular configuration to the ventricular, pulmonary artery, or peripheral arteries, while their disadvantage is weak gripping. Basket catheters have a powerful grasp and are capable of withdrawing relatively large foreign bodies. The diameter of the basket can be adjusted according to the vessel diameter. This catheter is preferred in situations where a foreign body is attached to the vessel wall without a free edge (21). Grasping forceps can reach into small vessels, and their gripping power is advantageous in removing a foreign body strongly adhering to the vessel wall; however, controlling the catheter head is difficult (22). Nowadays, transcatheter retrieval is used in more than 90% of cases of removal of foreign bodies (21, 23-25).

The specific type of retrieval catheter used is determined by the size of the patients, the type of foreign body, and

exactly how and where the foreign body is situated within the vascular system. Then, directly through the sheath or through a catheter delivered through the sheath, the particular retrieval device is advanced to the foreign body and manipulated to grasp it. Once firmly grasped, the foreign body material is withdrawn into the large sheath and out of the body through the sheath. With the use of the large, long sheaths with these retrieval devices, it is usually no longer necessary to perform a venous cut down even for the final removal of the foreign body from the vessel or skin entry site (26).

Advanced devices have improved the success rate of foreign body retrieval, however when percutaneous intravascular removal is not successful for example when the foreign bodies are large and risk of severe complication during retrieval (e.g., arrhythmias, perforation of vessel wall and ventricles, artery spasm, thromboembolism, or vessel damage) is high, extracorporeal circulation is used for retrieval (27). Large fragments should be repositioned to the femoral vein and removed by surgical cut-down (28).

Unfortunately, because of the small size of the child, our used sheath was not large enough for the migrated catheter and it could not be retrieved through the sheath. Thus, we pulled it into the femoral vein and then removed it by performing a cut-down.

In conclusion a rare complication in the use of central catheters is fracture and cardiac embolization. These can be retrieved safely using a percutaneous snare-catheter. We offer gentle bringing out of the catheter lines under the guide of fluoroscopy in all of the cases, if this is technically possible and safe.

Acknowledgments

The authors would like to thank the personnel of Fatemeh Zahra Cardiac Center for their cooperation and also to Miss Fatemeh Hosseinzadeh for editing the manuscript.

Conflict of Interest: None declared.

References

1. Lanzkowsky P. Manual of pediatric hematology and oncology, supportive care and management of oncology emergencies. 4 th ed. Chennai, India: Mcmilan Co 2007: 695-748.
2. Kliegman RM, Behrman RE, Jenson HB, et al. Nelson textbook of pediatrics. 18 th ed. Philadelphia, PA: Saunders Elsevier 2007; p: 383.
3. Ruiz CE, Nystrom GA, Butt AL, Zhang HP. Percutaneous retrieval of a broken umbilical catheter from left atrium in a remature newborn. *Cathet Cardiovasc Diagn* 1995; 36: 265-8.
4. Massin M, Lombet J, Rigo J. Percutaneous retrieval of broken silastic catheter from the left atrium in a critically III premature infant. *Cathet Cardiovasc Diagn* 1997; 42: 409-11.
5. Young T, Shaddy RE. Successful percutaneous retrieval of an intracardiac catheter fragment in an 800-g premature infant. *J Perinatol* 2003; 23: 67-8.
6. Simon-Fayard EE, Kroncke RS, Solarte D, Peverini R. Nonsurgical retrieval of embolized umbilical catheters in premature infants. *J Perinatol* 1997; 17: 143-7.
7. Schummer W, Schummer C, Gaser E, Bartunek R. Loss of the guide wire: mishap or blunder? *Br J Anaesth* 2002; 88: 144-6.
8. Kusminsky RE. Complications of central venous catheterization. *J Am Coll Surg* 2007; 204: 681-96.
9. Furutani S. Experience of a case with accidental retention of an IVH catheter in a blood vessel. *Okayama Red Cross Hosp J Med* 2005; 16: 25-30. [In Japanese]
10. Sadaoka S, Narita K, Fukuda K. Central venous catheter fragment with embolization into a distal branch of the pulmonary artery in a baby. *Jpn J Interv Radiol* 2004; 19: 237. [In Japanese]
11. Richardson JD, Grover FL, Trinkle JK. Intravenous catheter emboli. Experience with twenty cases and collective review. *Am J Surg* 1974; 128: 722-7.
12. Turner DD, Sommers SC. Accidental passage of a polyethylene catheter from cubital vein to right atrium: report of a fatal case. *N Engl J Med* 1954; 251: 744-5.
13. Cope C. Intravascular breakage of Seldinger spring guide wires. *JAMA* 1962; 180: 1061-3.
14. Seldinger SI. Catheter replacement of the needle in percutaneous arteriography: a new technique. *Acta Radiol* 1953; 39: 368-76.
15. Thomas J, Sinclair-Smith B, Bloomfield D, Davachi A. Nonsurgical retrieval of a broken segment of steel spring guide from the right atrium and inferior vena cava. *Circulation* 1964; 30: 106-8.
16. Dotter CT, Rosch J, Bilbao MK. Transluminal extraction of catheter and guide fragments from the heart and great

- vessels: 29 collected cases. *Am J Roentgenol Radium Ther Nucl Med* 1971; 111: 467–72.
17. Uflacker R, Lima S, Melichar AC. Intravascular foreign bodies: percutaneous retrieval. *Radiology* 1986; 160: 731–5.
 18. Cekirge S, Weiss JP, Foster RG, Neiman HL, McLean GK. Percutaneous retrieval of foreign bodies: experience with the nitinol goose neck snare. *J Vasc Interv Radiol* 1993; 4: 805–10.
 19. Curry JL. Recovery of detached intravascular catheter or guide wire fragments. A proposed method. *Am J Roentgenol Radium Ther Nucl Med* 1969; 105: 894–6.
 20. Sano A. Percutaneous retrieval of endovascular foreign bodies – experiences of Amplatz goose-neck snare. *Surg Ther* 1995; 73: 173–9. [In Japanese]
 21. Saitoh T, Ohtaki M, Matusyama M. Percutaneous intravascular removal of foreign body. *Medicina* 1996; 33: 1086–9. [In Japanese]
 22. Kuroki K. Intravascular removal of foreign body (from the right heart). In: Kumazaki T, Nakajima Y (eds) *IVR of emergency cases*. Tokyo: Medical View 2003; pp: 130–1.
 23. Savage C, Ozkan OS, Walser EM, Wang D, Zwischenberger JB. Percutaneous retrieval of chronic intravascular foreign bodies. *Cardiovasc Intervent Radiol* 2003; 26: 440–2.
 24. Yedlicka JW Jr, Carlson JE, Hunter DW, Castaneda-Zuniga WR, Amplatz K. Nitinol gooseneck snare for removal of foreign bodies: experimental study and clinical evaluation. *Radiology* 1991; 178: 691–3.
 25. Gabelmann A, Kramer S, Gorich J. Percutaneous retrieval of lost or misplaced intravascular objects. *AJR Am J Roentgenol* 2001; 176: 1509–13.
 26. Holzer RJ, Cheatham JP. Therapeutic method, therapeutic cardiac catheterization. In: Allen HD, Driscoll DJ, Shaddy RE, Feltes TF. *Moss and Adam's heart disease in infants, children and adolescents including the fetus and young adult*. 7 th ed. Philadelphia, PA: Lippincott Williams & wilkins 2008; p: 391.
 27. Actis Dato GM, Aidala E, Zattera GF, LeMaire SA, Wall MJ J., Mattox KL. Foreign bodies in the heart: surgical or medical therapy? *Ann Thorac Surg* 1999; 68: 291–2.
 28. Egglin TKP, Dickey KW, Rosenblatt M, Pollak JS. Retrieval of intravascular foreign bodies: experience in 32 cases. *AJR Am J Roentgenol* 1995; 164: 1259–64.