

## Short Communication

Masoud Pezeshki Rad (MD) <sup>1</sup>  
 Bita Abbasi (MD) <sup>1</sup>  
 Niloufar Valizadeh (MD) <sup>2</sup>  
 Farbod hatami (MD) <sup>2</sup>  
 Fariba tohidinezhad (PhD) <sup>3</sup>  
 Zahra Gharehbaghi (MD) <sup>1\*</sup>

1. Department of Radiology,  
 Mashhad University of Medical  
 Sciences, Mashhad, Iran

2. Cardiovascular Diseases  
 Research Center, Birjand  
 University of Medical Sciences,  
 Birjand, Iran

3. Department of Medical  
 Informatics, Mashhad University of  
 Medical Sciences, Mashhad, Iran

## \* Correspondence:

Zahra Gharehbaghi, Department  
 of Radiology, Mashhad University  
 of Medical Sciences, Mashhad, Iran

## E-mail:

zahrgharehbaghi6969@gmail.com

Tel: 0098 3841 1538

Fax: 0098 38430249

Received: 6 May 2021

Revised: 3 July 2021

Accepted: 28 Aug 2021

## Evaluation of Normal Renal Size and its Influencing Factors: A Cross-Sectional Study on the Adult Population of Mashhad

### Abstract

**Background:** The normal range of kidney size is a controversial issue among different populations given to its impressibility by multiple factors, therefore, this study aimed to provide valid reference ranges for kidney dimensions in the adult population of Mashhad. Also, we assessed the association of kidney size characteristics with some personal predisposing factors.

**Methods:** This cross-sectional study was conducted on 938 healthy individuals. Ultrasound measurement, physical examination, and laboratory analysis were performed. Demographic, dietary, and anthropometric data were obtained. The variables were categorized into 5 groups each, and data analysis were performed using the following statistical tests: Pearson correlation test, variance analysis, t-test, and chi-square test. A value of  $p < 0.05$  was considered statistically significant.

**Results:** Weight had the most association with kidney size followed to a lesser extent by height and age. Even after adjustment for other confounding variables, weight remained as an independent factor, while this effect was resolved for height and age. Also, all values for renal function, body bio-impedance, blood pressure components, and water intake were notably correlated with kidney size.

**Conclusion:** This study determined the normal kidney size in healthy adults. We also declared the normal range of kidney size is a dynamic concept and should be assessed for each individual separately according to their personal determinative factors.

**Keywords:** Kidney size, Ultrasound, Anthropometric characteristics, Renal function

### Citation:

Pezeshki Rad M, Abbasi B, Valizadeh n, et al. Evaluation of Normal Renal Size and its Influencing Factors: A Cross-Sectional Study on the Adult Population of Mashhad. Caspian J Intern Med 2022; 13(3): 623-633.

**R**enal size is an important and useful parameter for diagnosis, decision making, and clinical management of renal disorders regarding its close relationship to kidney function (1). The optimal ranges for the healthy adults' kidney size have been largely expressed by previous reports; herein, approximate estimations of each of the following parameters are 2.5-3 centimeters (cm) for thickness, 90-130 cm for length, and 5-5.7cm for width (2,3). Studies have shown remarkable differences in estimated normal kidney size between populations, which has been attributed to ethnic and racial diversity (4). Timely diagnosis and treatment of chronic kidney disease which is mainly known by a decrease in renal size can prevent the disease aggravation and its adverse consequents (5). Moreover, regarding anatomic landmarks, kidney size parameters are precisely required when performing a renal biopsy under ultrasound guidance. In the past several decades, ultrasonography has played an important role in evaluating renal size; as it is an inexpensive, accurate, non-invasive, radiation-free and readily available modality to determine length, width, and thickness of the organ (6).



Although, ultrasound is a well-established diagnostic method, yet there are insufficient data on the aspects of normal kidney size and its influencing factors in the healthy adult population. To the best of our knowledge, this is the first study in our region that aimed to evaluate the issue on a large sample volume.

## Methods

After approval by the local Ethics Committee from the Mashhad University of Medical Sciences Review Board (IR.MUMS.MEDICAL.RE.1397.104), this cross-sectional study was conducted at the radiology department of Imam Reza hospital in Mashhad during 2018-2019. Samples were selected by the available-sampling method and informed consent was obtained from all participants. Inclusion criteria were: age more than 18 years, serum creatinine  $\leq 1.5$  milligrams/deciliter (mg/dL), fasting blood sugar  $< 110$  mg/dL, and normal kidney appearance at the ultrasound.

The exclusion criteria included chronic or acute kidney disease, situations affecting kidney condition such as diabetes, hypertension, pregnancy, abnormalities in kidney anatomy such as renal cyst, and the patients who did not complete the study for any reason.

A total of 1000 individuals volunteered for the current study while 62 patients were excluded, the staff of Mashhad University of Medical Sciences who voluntarily participated were examined. The administrative unit made contact with the cases and the individuals selected their desired time and date and after registering the appointment, they were referred to the research unit. Each person received an identification code before the investigation.

Thirteen stations were defined for data collection. For all the participants, body composition analysis was performed via the bioimpedance method, by Bioelectrical Impedance Analysis Equipment, which evaluates the amount of body protein, intracellular and extracellular water, and minerals based on the absorption of electromagnetic rays. Likewise, serum creatinine and urea levels were evaluated, and also anthropometric characteristics including height, weight, body mass index (BMI), body surface area (BSA), and abdominal circumference were recorded. Each individual underwent renal ultrasonography to measure kidney length, width, thickness, and *anteroposterior* diameter. The procedure was performed using the Philips Affinity 50G device with a C6-2 Convex probe and a frequency of 3.2 HZ in longitudinal and

transverse view point in the supine position. Parenchymal thickness was measured in three (upper, middle, and lower) poles for each individual (figure 1).

The results were collected in a pre-designed checklist. Data analysis was performed via SPSS software Version 21 (SPSS Inc., Chicago, IL, USA) using statistical methods including descriptive and inferential statistics. Data were presented as the mean  $\pm$  standard deviation and the t-test, chi-square, and variance tests were used to identify relationships. Bivariate and multivariate analysis was performed to examine the effect of possible confounders. A value of  $P < 0.05$  was considered statistically significant. The datasets supporting the conclusions of this article are available in the radiology department of Imam Reza Hospital data base (Mashhad, Iran) repository. Furthermore, the datasets analyzed during the current study are available from the corresponding author on reasonable request.

## Results

For the present study, we analyzed the data collected from 938 individuals, of which 444 were males and 494 were females. The average age of men was  $46 \pm 8.1$  years and for women was  $43.4 \pm 6.9$ . Our cases were mostly Persian, while no significant difference was noted among different ethnicities including Persian, Turk, Kurd, and others regarding gender. While all the anthropometric, bio-impedance, biochemistry, blood pressure, and ultrasound measures mentioned were higher in the male group ( $p < 0.001$ ), BMI and dietary intakes did not have a notable difference between two gender groups (table 1).

Findings from ultrasonic measurements revealed significantly greater values for Right Renal Length (RRL), Left Renal Length (LRL), Right Renal Width (RRW), and Left Renal Width (LRW) in men compared to women ( $p < 0.001$ ). A significant difference in renal size was noted in female gender when considering age groups, so, the age group of 41-50 years had an increase in both right ( $104.7 \pm 8.4$  mm,  $p = 0.004$ ) and left ( $104.9 \pm 10.8$  mm,  $P = 0.039$ ) kidney length in comparison to other age groups (table 2).

In both kidneys, the mean values for parenchymal thickness and renal length were statistically higher in men and women over 180 cm and 170 cm of height, respectively (table 3). Our results demonstrated that in men weighing over 100 kg, an increase in all parameters of kidney size was observable ( $p < 0.05$ ). In the women group, tests showed a significant

association between the weight and all kidney size parameters except for Left Renal Parenchymal Thickness (LRP) ( $p < 0.05$ ). The mean width and parenchymal thickness of both kidneys were notably higher than other weight groups in women weighing 90-100 kg and over 100 kg respectively (table 4). We performed the Pearson correlation test to assess the role of possible factors contributing to the variation of the kidney sizes. Results of the Pearson test confirmed a significant positive correlation between all the variables of the current study including age, anthropometric characteristics, body composition components, blood pressure values, laboratory values, amount of water intake, and kidney size

parameters. This effect was not seen in the case of salt intake. Details are presented in table 5. Based on the findings of the single-variable regression test (table 6), the relationship between height, weight, abdominal circumference, body water, minerals, protein, systolic, and diastolic blood pressure and Mean Arterial Pressure (MAP) with kidney length was significant. Also, based on the findings of multiple regressions analysis, the relationships of 3 variables (weight, height, and minerals) with kidney length were significant (figures 2, 3). The value of the determination coefficient for the final model was 0.27 ( $\beta = 97.1$ ), which indicates the prediction of 27% of changes in kidney thickness by these variables.

**Table 1. Basic demographic and clinical characteristics of the population (N=938)**

Variable	Male (N=444)	Female (N=494)	P-value
Age (year)	46.0 ± 8.1	43.4 ± 6.9	<0.001
Anthropometric			
Height (cm)	172.9 ± 6.9	158.3 ± 5.9	<0.001
Weight (kg)	80.6 ± 12.5	68.3 ± 11.3	<0.001
BMI (kg/m <sup>2</sup> )	26.9 ± 3.6	27.3 ± 4.3	0.185
BSA (m <sup>2</sup> )	1.9 ± 0.2	1.7 ± 0.1	<0.001
Abdomen Circumference(cm)	96.0 ± 10.6	91.8 ± 10.2	<0.001
Bio-impedance			
Body Water (L)			
Total	42.4 ± 5.6	29.8 ± 3.6	<0.001
ICW	26.4 ± 3.5	18.4 ± 2.2	<0.001
ECW	16.0 ± 2.1	11.5 ± 1.4	<0.001
Mineral (kg)	3.9 ± 0.6	2.8 ± 0.3	<0.001
Protein (kg)	11.4 ± 1.5	7.9 ± 1.0	<0.001
Biochemistry			
BUN	32.8 ± 6.4	28.9 ± 6.9	<0.001
Creatinine	1.3 ± 0.2	1.1 ± 0.2	<0.001
Blood Pressure (mmHg)			
Systolic	110.3 ± 22.1	99.0 ± 18.4	<0.001
Diastolic	70.7 ± 14.6	64.9 ± 11.5	<0.001
MAP	97.9 ± 12.3	89.6 ± 11.7	<0.001
Dietary Intake			
Water (glass per day)	5.0 ± 3.0	4.6 ± 3.3	0.103
Salt (gram per day)	2.8 ± 3.0	3.1 ± 3.7	0.186
Ultrasound Measures (mm)			
RRL	107.6 ± 8.6	103.8 ± 8.3	<0.001
RRW	43.7 ± 7.4	38.2 ± 6.0	<0.001
LRL	108.9 ± 7.9	104.1 ± 10.0	<0.001
LRW	47.9 ± 7.1	43.8 ± 6.8	<0.001

BMI, Body Mass Index; BSA, Body Surface Area; ICW, Intracellular Water; ECW, Extracellular Water; BUN, Blood Urea Nitrogen; MAP, Mean Arterial Pressure; RRL=Right Renal Length, LRL=Left Renal Length, RRW=Right Renal Width, LRW=Left Renal Width

**Notes:** Values are represented as the mean ± SD or N (%). BSA was calculated using the DuBois formula. Statistical analysis via independent - test or chi-square.

**Table 2: Normal values of the kidney measurements classified Bbsed on age and gender category**

Gender	Measures (mm)	30-40 (N= 175)	41-50 (N=179)	51-60 (N=83)	61-70 (N=7)	P-value
Male	RRL	108.6±8.1 (83-130)	106.5±8.9 (86-130)	106.6±7.6 (90-132)	102.6±7.3 (95-112)	0.062
	RRE	43.9±7.5 (31-91)	42.7±6.6 (26-65)	43.1±8.4 (28-99)	43.3±6.9 (30-53)	0.671
	RR APD	46.1±6.6 (29-71)	45.6±5.9 (32-62)	45.5±5.7 (35-60)	47.1±4.8 (42-52)	0.815
	RRP	14.9±2.8 (9.7-26)	14.8±2.6 (9.3-22)	14.6±2.7 (9-23.7)	14.3±2.6 (10-8.3)	0.695
	LRL	109.0±7.7 (86-133)	109.1±7.6 (94-132)	107.6±7.2 (94-129)	102.0±9.7 (86-114)	0.062
	LRW	48.1±6.7 (30-64)	47.4±6.7 (30-66)	46.5±7.4 (10-63)	45.9±5.4 (38-55)	0.360
	LR APD	46.9±5.8 (33-64)	47.0±7.1 (18-67)	46.9±6.3 (28-64.5)	45.9±7.3 (31-54)	0.973
	LRP	15.7±3.0 (8.7-28)	15.4±2.8 (9-29)	15.4±2.7 (9-24.7)	15.0±3.1 (10-1.3)	0.706
		30-40 (N=235)	41-50 (N=193)	51-60 (N=56)	61-70 (N=10)	P-value
	Female	RRL	104.3±8.3 (84-126)	104.7±8.4 (85-130)	100.8±7.2 (85-117)	99.2±7.8 (85-108)
RRW		37.8±5.8 (22-56)	38.5±6.2 (25-63)	37.7±5.8 (27-53)	39.6±7.5 (23-47)	0.570
RR APD		42.7±6.2 (17-60)	42.9±6.2 (28-65)	43.8±5.5 (32-59)	43.6±6.1 (35-55)	0.712
RRP		13.8±2.7 (9-24)	13.9±2.6 (6.3-21.3)	13.2±2.3 (8.3-18.5)	13.8±3.1 (10-19.5)	0.219
LRL		104.3±0.4 (11-132)	104.9±10.8 (10-123)	102.0±0.1 (88-117)	96.6±6.7 (90-111)	<b>0.039</b>
LRW		44.1±6.9 (13-58)	42.9±7.0 (11-59)	44.8±5.9 (28-57)	40.7±7.3 (27-50)	0.095
LR APD		44.5±5.2 (33-59)	44.3±6.1 (7-62)	44.1±5.6 (30-58)	42.7±7.1 (30-56)	0.757
LRP		14.5±2.7 (7.3-24.7)	14.7±2.7 (8.7-23.3)	14.1±2.5 (8.3-19.7)	14.5±.9 (9.7-8.3)	0.417

RRL=Right Renal Length, LRL=Left Renal Length, RRW=Right Renal Width, LRW=Left Renal Width, RRT=Right Renal Parenchymal Thickness, LRP=Left Renal Parenchymal Thickness, RR APD=Right Renal Anteroposterior Diameter, LR APD=Left Renal Anteroposterior Diameter;

Notes: Values are represented as mean ± SD (min-max).

**Table 3. Normal values of the kidney measurements classified based on the height and gender category**

Gender	Measures (mm)	140-150 (N=0)	151-160 (N=13)	161-170 (N=131)	171-180 (N=253)	>180 (N=47)	P-value
Male	RRL	N/A	102 ± 6.6 (90-110)	106 ± 8.1 (86-130)	107 ± 8.2 (83-130)	112 ± 7.9 (97-132)	<b>&lt;0.001</b>
	RRW	N/A	41 ± 7.5 (30-53)	43 ± 5.6 (30-63)	43 ± 8.4 (26-99)	46 ± 6.4 (32-58)	0.073

	RR APD	N/A	44 ± 7.0 (29-56)	45 ± 6.0 (32-67)	46 ± 6.0 (33-71)	48 ± 6.3 (36-62)	0.065	
	RRP	N/A	13.2 ± 2.0 (10-17.3)	14.7 ± 2.6 (9-26)	14.7 ± 2.6 (9.3-25.5)	15.7 ± 3.0 (10-23.7)	<b>0.004</b>	
	LRL	N/A	102 ± 9.1 (86-116)	108 ± 7.1 (93-125)	109 ± 7.8 (86-133)	111 ± 6.7 (95-128)	<b>0.001</b>	
	LRW	N/A	46 ± 5.5 (38-55)	47 ± 7.5 (10-61)	47 ± 6.4 (30-66)	50 ± 7.0 (30-64)	0.086	
	LR APD	N/A	46 ± 4.2 (41-53)	46 ± 5.7 (31-65)	47 ± 6.9 (18-67)	49 ± 6.2 (35-64)	0.109	
	LRP	N/A	13.7 ± 2.4 (9-17)	15.5 ± 2.9 (8.7-28)	15.4 ± 2.7 (9-29)	16.2 ± 2.9 (9.7-25.2)	<b>0.009</b>	
			<b>140-150 (N=31)</b>	<b>151-160 (N=271)</b>	<b>161-170 (N=183)</b>	<b>171-180 (N=9)</b>	<b>&gt;180 (N=0)</b>	<b>P-value</b>
<b>Female</b>	RRL		100 ± 7.9 (85-120)	103 ± 7.7 (84-130)	106 ± 8.6 (86-128)	109 ± 5.6 (100-114)	N/A	<b>&lt;0.001</b>
	RRW		36 ± 5.2 (23-49)	38 ± 6.2 (22-61)	38 ± 5.8 (25-63)	39 ± 6.5 (30-46)	N/A	0.083
	RR APD		40 ± 4.1 (32-50)	43 ± 5.9 (30-60)	43 ± 6.5 (17-65)	41 ± 6.9 (34-51)	N/A	0.067
	RRP		12.9±2.7 (8.3-19.7)	13.7 ± 2.5 (6.3-21.7)	13.9 ± 2.8 (7.3-24)	14.7 ± 2.7 (10.3-17.5)	N/A	0.226
	LRL		96±17.1 (10-113)	104 ± 7.5 (83-123)	106 ± 11.2 (11-132)	111 ± 6.1 (103-120)	N/A	<b>&lt;0.001</b>
	LRW		43 ± 7.1 (27-56)	44 ± 7.0 (11-59)	44 ± 6.6 (16-57)	45 ± 3.1 (42-50)	N/A	0.951
	LR APD		42 ± 4.8 (30-54)	44 ± 5.5 (30-59)	45 ± 5.9 (7-62)	46 ± 3.1 (42-49)	N/A	0.158
	LRP		14.2±2.4 (9.3-20)	14.5 ± 2.5 (7.3-22)	14.6 ± 2.9 (8-24.7)	16.1 ± 2.3 (12.7-19.3)	N/A	0.513

RRL=Right Renal Length, LRL=Left Renal Length, RRW=Right Renal Width, LRW=Left Renal Width, RRT=Right Renal Parenchymal Thickness, LRP=Left Renal Parenchymal Thickness, RR APD=Right Renal Anteroposterior Diameter, LR APD=Left Renal Anteroposterior Diameter;

Notes: Values are represented as mean ± SD (min-max).

**Table 4. Normal values of the kidney measurements classified based on the gender and weight category**

Gender	Measures (mm)	<60 * (N=11)	60-70 (N=85)	71-80 (N=130)	81-90 (N=145)	91-100 (N=55)	>100 (N=18)	P-value
<b>Male</b>	RRL	98 ± 8.0 (86-112)	104 ± 7.3 (83-117)	106 ± 7.1 (87-125)	109 ± 7.5 (92-130)	112 ± 8.4 (92-132)	115 ± 9.6 (93-128)	<b>&lt;0.001</b>
	RRW	37 ± 6.4 (30-50)	40 ± 5.6 (26-53)	42 ± 8.8 (30-99)	45 ± 5.6 (34-65)	47 ± 6.3 (34-65)	48 ± 5.3 (38-58)	<b>&lt;0.001</b>
	RR APD	39 ± 6.0 (29-48)	44 ± 4.9 (36-55)	45 ± 6.1 (33-71)	46 ± 5.8 (35-67)	49 ± 6.3 (39-64)	50 ± 6.0 (40-60)	<b>&lt;0.001</b>
	RRP	13.6 ± 2.4 (9.3-18)	14.0 ± 2.3 (9.7-20)	14.2 ± 2.5 (9.7-20.3)	15.3 ± 2.7 (9-26)	15.2 ± 3.0 (10-25.5)	16.7 ± 3.0 (12-23.7)	<b>&lt;0.001</b>

	LRL	99 ± 5.6 (93-112)	106 ± 6.7 (92-119)	107 ± 6.6 (86-124)	110 ± 7.0 (95-130)	113 ± 8.6 (86-133)	117 ± 5.9 (108-129)	<0.001
	LRW	43 ± 4.3 (38-52)	44 ± 7.9 (10-58)	46 ± 6.4 (30-63)	49 ± 5.7 (32-65)	51 ± 6.7 (35-66)	49 ± 4.7 (41-56)	<0.001
	LR APD	42 ± 5.7 (31-53)	45 ± 6.2 (28-65)	46 ± 6.3 (18-59)	48 ± 6.1 (33-67)	50 ± 6.3 (38-65)	50 ± 6.0 (40-56)	<0.001
	LRP	14.1 ± 2.5 (9-18.5)	14.9 ± 2.7 (8.7-24.7)	15.2 ± 3.0 (9-29)	15.7 ± 2.6 (9.7-28)	15.9 ± 3.1 (10.7-25.2)	17.0 ± 2.7 (12-20.3)	0.005
		<60 (N=116)	60-70 (N=151)	71-80 (N=148)	81-90 (N=58)	91-100 (N=17)	>100 (N=4)	P- value
<b>Female</b>	RRL	101 ± 6.5 (86-118)	103 ± 7.8 (84-125)	106 ± 7.9 (89-124)	110 ± 8.5 (91-128)	109 ± 11.3 (85-130)	115 ± 12.6 (103-128)	<0.001
	RRW	36 ± 5.0 (22-53)	37 ± 5.7 (23-56)	40 ± 5.2 (27-53)	42 ± 7.1 (25-63)	46 ± 6.3 (36-61)	39 ± 6.5 (34-48)	<0.001
	RR APD	42 ± 5.8 (30-57)	42 ± 5.8 (28-59)	44 ± 6.5 (17-59)	44 ± 5.2 (34-58)	46 ± 5.5 (40-60)	50 ± 10.5 (42-65)	0.002
	RRP	10.6 ± 0.4 (10.3-10.8)	13.0 ± 2.6 (8.3-16.8)	13.7 ± 2.5 (9.3-21.2)	13.3 ± 2.5 (6.3-21)	14.2 ± 2.7 (8.3-24)	15.0 ± 3.4 (11.7-19)	0.018
	LRL	100 ± 11.1 (10-119)	104 ± 10.8 (11-123)	106 ± 7.2 (87-121)	108 ± 7.3 (93-122)	110 ± 10.9 (90-132)	110 ± 7.4 (100-117)	<0.001
	LRW	43 ± 7.1 (13-57)	43 ± 7.0 (11-59)	45 ± 6.3 (30-58)	46 ± 4.8 (38-57)	46 ± 8.0 (30-54)	42 ± 6.6 (34-48)	0.013
	LR APD	43 ± 5.1 (30-58)	44 ± 6.0 (7-59)	45 ± 5.6 (31-59)	46 ± 4.7 (32-57)	46 ± 4.6 (38-54)	47 ± 10.3 (40-62)	0.047
	LRP	12.2 ± 4.0 (9.3-15)	13.8 ± 1.7 (11.7-17)	14.4 ± 2.5 (7.3-20)	14.4 ± 2.6 (9-23.3)	14.8 ± 2.9 (8.3-24.7)	15.3 ± 2.3 (13.3-17.3)	0.374

RRL=Right Renal Length, LRL=Left Renal Length, RRW=Right Renal Width, LRW=Left Renal Width, RRT=Right Renal Parenchymal Thickness, LRP=Left Renal Parenchymal Thickness, RR APD=Right Renal Anteroposterior Diameter, LR APD=Left Renal Anteroposterior Diameter; \* kilogram

Notes: Values are represented as mean ± SD.

**Table 5: Correlation of the potential risk factors with kidney measures**

	RK length	RK width	RK AP	LK length	LK width	LK AP
Age	r=-0.040, P=0.219	r=0.095**, P=0.004	r=0.040, P=0.224	r=-0.024, P=0.467	r=0.021, P=0.528	r=0.018, P=0.585
Height	r=0.315**, P=0.000	r=0.339**, P=0.000	r=0.205**, P=0.000	r=0.305**, P=0.000	r=0.240**, P=0.000	r=0.220**, P=0.000
Weight	r=0.448**, P=0.000	r=0.450**, P=0.000	r=0.310**, P=0.000	r=0.415**, P=0.000	r=0.336**, P=0.000	r=0.302**, P=0.000
BMI	r=0.291**, P=0.000	r=0.275**, P=0.000	r=0.211**, P=0.000	r=0.258**, P=0.000	r=0.213**, P=0.000	r=0.191**, P=0.000
BSA	r=0.445**, P=0.000	r=0.457**, P=0.000	r=0.303**, P=0.000	r=0.418**, P=0.000	r=0.336**, P=0.000	r=0.304**, P=0.000
Abdomen Circumference	r=0.358**, P=0.000	r=0.369**, P=0.000	r=0.262**, P=0.000	r=0.330**, P=0.000	r=0.266**, P=0.000	r=0.241**, P=0.000

Total Body Water	r=0.407**, P=0.000	r=0.447**, P=0.000	r=0.288**, P=0.000	r=0.399**, P=0.000	r=0.343**, P=0.000	r=0.287**, P=0.000
ICW	r=0.406**, P=0.000	r=0.446**, P=0.000	r=0.287**, P=0.000	r=0.398**, P=0.000	r=0.344**, P=0.000	r=0.286**, P=0.000
ECW	r=0.408**, P=0.000	r=0.448**, P=0.000	r=0.289**, P=0.000	r=0.399**, P=0.000	r=0.338**, P=0.000	r=0.288**, P=0.000
Mineral (kg)	r=0.410**, P=0.000	r=0.431**, P=0.000	r=0.281**, P=0.000	r=0.394**, P=0.000	r=0.330**, P=0.000	r=0.282**, P=0.000
Protein (kg)	r=0.406**, P=0.000	r=0.446**, P=0.000	r=0.286**, P=0.000	r=0.399**, P=0.000	r=0.343**, P=0.000	r=0.285**, P=0.000
BUN	r=-0.033, P=0.330	r=0.058, P=0.083	r=0.049, P=0.150	r=0.003, P=0.928	r=0.067*, P=0.048	r=0.052, P=0.123
Creatinine	r=0.014, P=0.667	r=0.194**, P=0.000	r=0.112**, P=0.001	r=0.072*, P=0.032	r=0.106**, P=0.002	r=0.070*, P=0.037
Systolic	r=0.182**, P=0.000	r=0.155**, P=0.000	r=0.134**, P=0.000	r=0.165**, P=0.000	r=0.137**, P=0.000	r=0.097**, P=0.003
Diastolic	r=0.164**, P=0.000	r=0.151**, P=0.000	r=0.102**, P=0.002	r=0.139**, P=0.000	r=0.109**, P=0.001	r=0.076*, P=0.021
MAP	r=0.165**, P=0.000	r=0.184**, P=0.000	r=0.123**, P=0.000	r=0.186**, P=0.000	r=0.171**, P=0.000	r=0.081*, P=0.015
Water Intake	r=0.099**, P=.002	r=0.085**, P=.009	r=0.041, P=0.209	r=0.078*, P=0.017	r=0.134**, P=0.000	r=0.093**, P=0.005
Salt Intake	r=0.048, P=0.145	r=0.019, P=0.567	r=0.036, P=0.273	r=0.001, P=0.967	r=0.013, P=0.698	r=0.024, P=0.458


BMI=Body Mass Index; BSA=Body Surface Area; ICW=Intracellular Water; ECW= Extracellular Water; BUN=Blood Urea Nitrogen; MAP=Mean Arterial Pressure; RK, Right Kidney; LK, Left Kidney **Notes:** \*\*Correlation is significant at the 0.01 level (2-tailed). \*Correlation is significant at the 0.05 level (2-tailed).

**Table 6. Predictors of the kidney length (mean length of the right and left kidneys)**


Variable	Bivariate Analyses		Multivariate Analyses	
	$\beta$ (SE)	P	$\beta$ (SE)	P
Age (year)	-0.036 (0.034)	0.287		
Height	0.288 (0.025)	<0.001	-0.106 (0.057)	0.06
Weight	0.290 (0.017)	<0.001	0.189 (0.029)	<0.001
Abdomen Circumference (cm)	0.290 (0.022)	<0.001		
Total Body Water (L)	0.463 (0.029)	<0.001		
Mineral (kg)	5.094 (0.323)	<0.001	3.703 (0.989)	<0.001
Protein (kg)	1.689 (0.107)	<0.001		
BUN	-0.020 (0.038)	0.601		
Creatinine	1.801 (1.201)	0.134		
Systolic BP	0.074 (0.012)	<0.001		
Diastolic BP	0.101 (0.019)	<0.001		
MAP	0.125 (0.020)	<0.001		
Water Intake (glass per day)	0.255 (0.082)	0.002		
Salt Intake (gram per day)	0.056 (0.076)	0.460		

BUN, Blood Urea Nitrogen; MAP, Mean Arterial Pressure.

**Notes:** To adjust for multiple pairwise comparisons,  $P < .01$  is considered statistically significant. Using forward stepwise regression, 3 factors (height, weight, and mineral) were found to be associated with Kidney Width. The  $R^2$  value for the final model was 0.27 ( $\beta = 97.1$ ).




**Figure 1. Sonographic measurement of the kidney length and width and AP diameter**





**Figure 2. The Linear relationship of the height, weight, and minerals with right kidney sizes (length, width, and AP diameter)**



**Figure 3. The linear relationship of the height, weight, and minerals with left kidney sizes (length, width, and AP diameter)**

## Discussion

This study aimed to evaluate the normal renal size in the healthy Iranian adult population, and also to explore the possible correlation of the influencing factors on kidney size. There are discrepant reports about normal kidney size in different populations (4, 7). Nonetheless, few similar studies have been conducted to provide evidence for the standard kidney size in healthy populations and most of the studies have not excluded the conditions affecting renal size, such as diabetes and hypertension (8).

Although this variety is considered to be ethnic, we cannot only blame genetics. The environmental factors should also be considered as a significant factor (9). Therefore, it does not seem possible to determine an absolute number for normal kidney size. As mentioned earlier, measurement of kidney length is commonly used for diagnosis and treatment of kidney diseases and has a special place in the treatment, timely diagnosis, and preventing side effects (2). Our results showed

a significant difference in kidney size between men and women so, the male renal size was significantly larger (table1).As there are conflicting reports on the correlation of

kidney size with age (9-11) we also investigated this relationship. Our findings revealed aging and RRW had a significant positive correlation (table 5). When considering sex, the female population showed a higher RRL ( $p=0.004$ ) and LRL ( $P=0.039$ ) between the age of 41-50 years, while male kidney size was not significantly different regarding to the study's age groups (table 2). In a similar study by Jabbari et al., the analysis of different age groups, regardless of gender, showed a significant decrease in kidney length and parenchymal thickness after the age of fifty, according to their proportion of study population, which mostly included women, we have obtained the same results in our female population (9). In the study of Raza et al. in Pakistan, the relationship between the mean kidney length and age was found to be significant (11). Glodny et al. reported that kidney

length increased significantly until the fifth decade of life in men (10). According to differences between age groups of our study, the resulting discrepancy may be explained.

Based on our study, height had a significant and positive correlation with all renal measurements (table 5, figures 2,3), Just as many other studies with a large population reported (10-12), Also, some small studies described no correlation (8,13). As for sex differences, within the male population, length, and parenchymal thickness in both kidneys and in the female group, both renal lengths had a significant positive relation with body height. Since weight and BMI had a significant positive correlation with kidney size, regarding gender, weight was also related to all renal measurements (except female LRP), as many other authors had mentioned (8-12,14). In a study by Muthusami et al., a moderate positive correlation was reported between kidney length with body weight and BSA while they estimated a poor positive correlation with height and BMI (7).

According to our study, intracellular water, extracellular water, protein, and minerals also had a significant relationship with kidney measurement parameters. Furthermore, there was a positive and significant relationship between water consumption in the studied subjects and both kidneys' lengths and width and left anteroposterior diameter. However, no significant relationship was observed between the amount of salt diet and any of the mentioned variables.

To resolve the confounding effect of other variables, we performed the univariate and multivariate regression analysis. According to the studied variables, three variables including height, weight, and minerals remained significantly associated with kidney length (table 6). Su H-A et al. reported the association of the kidney length with weight and height (4). Also, Glondy et al. mentioned the same link for renal length and body height extracted from regression analysis (10).

The value of the determination coefficient for the final model was 0.27 ( $\beta = 97.1$ ), which indicates the prediction of 27% of changes in kidney thickness by these three variables. This study was conducted to assign reference ranges for normal kidney size in the Iranian adult population and to assess its possible influencing factors. The reference ranges presented in this study, as well as others, can be considered for implications in clinical guidelines or practice, however, in our opinion looking for an absolute normal kidney size definition is an unreasonable effort regarding its dynamic nature implying multiple factors capable to modify kidney size. We recommend researchers, clinicians, and radiologists

to define a normal kidney size for every person separately and individually concerning factors including sex, age, underlying diseases, anthropometric characteristics, and diet. Future studies are suggested to explore the role of such modifying factors with regard to race in larger populations with a long-term follow-up design.

### Acknowledgements

We appreciate very much Mashhad Persian Cohort Center and the personnel of Imam Reza Hospital for collaboration in this research.

**Funding:** This research received no specific grant from any funding agency.

**Conflicts of Interest:** The authors declare that there is no conflict of interest.

### Authors' contributions:

Concept/Design- M.P.R., B.A., Z.G. Acquisition of Data- M.P.R., B.A., Z.G., N.V. Data Analysis/Interpretation- M.P.R., B.A., Z.G., N.V., F.H., F.T. Drafting of the manuscript- Z.G., N.V., F.H. Critical revision of the manuscript and approval of the article- B.A., Z.G., N.V., F.H.

### References

1. Sutton D. Textbook of radiology and imaging. 7th ed. London: Churchill Livingstone 2003; pp: 885-988.
2. Rosenbaum DM, Korngold E, Teele RL. Sonographic assessment of renal length in normal children. *Am J Roentgenol* 1984; 142: 467-9.
3. Pollack HM, McClennan BL. Clinical urography. 2nd ed. Philadelphia: WB Saunders 2000; pp: 473-504.
4. Su HA, Hsieh HY, Lee CT, et al. Reference ranges for ultrasonographic renal dimensions as functions of age and body indices: A retrospective observational study in Taiwan. *Plos One* 2019; 14: e0224785.
5. Kasper DL, Braunwald E, Fauci AS. Harrison's principles of internal medicine. 16th ed. New York: McGraw Hill, 2005; pp: 1653-63.
6. Armstrong P, Wastie ML. Diagnostic imaging. 2nd ed. London: Blackwell 1990; pp: 203-348.
7. Muthusami P, Ananthakrishnan R, Santosh P. Need for a nomogram of renal sizes in the Indian population—findings from a single center sonographic study. *Indian J Med Res* 2014; 139: 686-93.

8. Okoye IJ, Agwu KK, Eze CU. Relationship between sonographic renal length and renal parenchymal thickness in normal adult southeast Nigerians. *West Afr J Med* 2006; 25: 231-4.
9. Jabbari M, Mollazade R, ESNA AF, Alizadeh Z. Normal renal dimensions in Iranian adults measured by ultrasound. *Anatomical Sci* 2016; 13: 25-32.
10. Glodny B, Unterholzner V, Taferner B, et al. Normal kidney size and its influencing factors-a 64-slice MDCT study of 1.040 asymptomatic patients. *BMC Urol* 2009; 9: 19.
11. Raza M, Hameed A, Khan MI. Ultrasonographic assessment of renal size and its correlation with body mass index in adults without known renal disease. *J Ayub Med Coll Abbottabad* 2011; 23: 64-8.
12. Kang KY, Lee YJ, Park SC, et al. A comparative study of methods of estimating kidney length in kidney transplantation donors. *Nephrol Dialysis Transplant* 2007; 22: 2322-7.
13. Barton EN, West WM, Sargeant LA, Lindo JF, Iheonunekwu NC. A sonographic study of kidney dimensions in a sample of healthy Jamaicans. *West Indian Med J* 2000; 49: 154-7.
14. El-Reshaid W, Abdul-Fattah H. Sonographic assessment of renal size in healthy adults. *Med Princ Pract* 2014; 23: 432-6.